



Laparoscopic Inguinal Hernia Repair Post-Op Instructions

Post-Op To-Do List:

- Take Acetaminophen and Ibuprofen tablets every 6 hours together for 2-5 days
- Do not do anything strenuous for 2 weeks after surgery
- Please review the post op instructions below if you have questions

Pain

Soreness or pain at the incisions, the lower abdomen, and groin after surgery is normal. Pain will be worse with movement, changing positions from laying to seated to standing. Male patients may notice new pain in the testicles after surgery. This is normal and will go away over time.

Most of our patients take only tylenol and ibuprofen for pain management after surgery. We recommend taking the following every 6 hours for the first few days after surgery as needed (both can be taken together at the same time):

- 650mg to 1g of acetaminophen (Tylenol)
- 400mg to 600mg of ibuprofen (Advil or Motrin)

If you are experiencing significant pain that is preventing you from going on a walk or falling asleep, please call the office for a prescription for oxycodone. This medication will cause constipation, so we recommend the following:

- Fiber supplement (Citrucel or Metamucil)
- Stool softener (Colace) or natural laxative (Senokot)
- Laxative only if unable to have a bowel movement for a few days (Milk of Magnesia)

Incision

The incisions are closed with dissolvable sutures - these do not need to be removed. You have glue over your incisions with dissolvable stitches under the skin. You should leave this glue in

place as long as it is well stuck in the area. If it peels up at the edges (usually 10-14 days after surgery), you may pull it off. Otherwise it will fall off on its own.

Your incisions may feel hard about 7-10 days after surgery. This is due to scar tissue, and is a normal part of healing.

Bruising in the groin and genitals is normal after surgery. MALE patients may find that your scrotum or penis will become discolored, and you will have swelling in the region of your hernia and incision. FEMALE patients may find that your labia will become swollen and discolored. This discoloration and swelling will resolve with time.

Some patients will feel a firm lump in the area where their hernia used to bulge. Most of the time this is fluid in this area - called a "seroma" - this will resolve within a few months.

Many patients have used Arnica Montana, either as a gel or orally, to reduce swelling and bruising after surgery. Arnica is a safe, homeopathic medicine that works well in many patients. Most natural food stores will carry it.

Bathing

You may shower at any time after surgery.

Please avoid baths, hot tubs, pools, or soaking in water for 2 weeks after surgery, as this can increase your risk of an infection.

Activity

Following surgery, you are encouraged to do as much walking as is comfortable. You may climb stairs, taking them one at a time and slowly. We recommend walking at a leisurely pace for the first 2 weeks.

Avoid exercise, strenuous activity, lifting over 25lbs, and repetitive actions for 2 weeks after surgery. After 2 weeks, you may slowly resume these activities, but do so slowly and progressively, and listen to your body if it is telling you to stop or slow down.

You may drive 24 hours after surgery if you feel up to it, as long as you are not taking any prescription pain medications. We typically recommend you avoid driving for 48-72 hours unless necessary.

Precautions

Although not commonly seen, any incision is susceptible to infection. If you develop a fever of 101 degrees or above, have unexpected pain, redness or pus draining from the incision, please contact our office.

Your pain and swelling should gradually improve after the second or third postoperative day. If you have new, constant pain after feeling good for a couple of days, please contact our office.

Follow-up

You will have a post-operative appointment scheduled for somewhere between 2 and 4 weeks after surgery. If this has not been scheduled yet, or if you need to reschedule, please contact our office.

Hernia recovery during cold and flu season

The winter months bring lots of colds and sometimes the flu. If you find yourself needing to cough or sneeze, it's often helpful to place a pillow or your hand over your incisions. This provides some support and comfort, and will hopefully decrease your pain when you cough or sneeze. Sharp pain is not uncommon after a cough or sneeze, and lingering effects should resolve in a day or two. It's very unlikely that the hernia will return as a result of this.

Post-op FAQ:

Below are questions that patients often ask us before and after surgery. This was created to help you better prepare for the day of surgery and for recovery. While these answers do not substitute for an appointment with us, they are created to help you remember what may have been said in the office during your consultation.

Will I be able to walk after surgery?

Yes. We want everyone to walk after surgery. Rarely, the local anesthesia we use can cause your leg to be numb for several hours. In this case, you will need to use crutches for a few hours after surgery.

When can I shower after surgery?

You may shower the day after surgery. There is waterproof glue over the incisions.

When should I take my dressing off?

You have glue over your incisions with dissolvable stitches under the skin. You should leave this glue in place as long as it is well stuck in the area. If it peels up at the edges (usually 10-14 days after surgery), you may pull it off. Otherwise it will fall off on its own.

Can I put a bandaid over the glue?

Sure. Some people find that a bandaid feels better. Sometimes a bandaid can help keep your clothes clean in the small chance a few drops of blood seep from under the glue.

When can I drive after surgery?

You can drive anytime after 24 hours post-op, unless you are taking opioids for pain. You CANNOT drive if taking the prescription opioids. We suggest waiting 24 full hours after your last dose of prescription pain medication before you drive. If you have not taken prescription medication or finished them, you may drive if you feel that you are able to.

Can I go up steps after my hernia surgery?

Yes. Please take steps slowly and one at a time.

Can I lift a gallon of milk?

Yes, as long as it's once or twice! You may lift up to 25 lbs as long as you are not performing many repetitive motions the first 2 weeks. Repetitive activities can cause harm to your repair.

Can I bend down to tie my shoes after surgery?

Yes, just move slowly.

Can I go for a long walk after hernia surgery?

Yes, we encourage it. Just avoid strenuous hiking trails and don't wear a backpack.

When can I go for a run after surgery?

You can go for a run 2 weeks after hernia surgery.

When can I resume sexual activity?

You may resume sexual activity 2 weeks after hernia surgery.

When can I go back to work after hernia surgery?

You may go to work whenever you feel like, as long as your job does not require heavy lifting. Approximately 75% of our patients go to work within 3 days of surgery.

When can I go back to the gym?

You may return to the gym 2 weeks after surgery. Please use common sense and resume activity gradually.

How much pain will I have after surgery?

Everyone is different! Some pain, of course, is expected. Most of our patients feel that Tylenol and Ibuprofen are strong enough to take care of their pain.

Is testicle pain normal after surgery?

Yes, some patients will have irritation or sensitivity of the nerves that go down to the testicle, therefore causing pain or sensitivity of the testicle. This is normal and will resolve with time.

How likely is the hernia to come back after surgery?

The chance that an inguinal hernia comes back after no mesh surgery is about 2%.
The chance that an inguinal hernia comes back after mesh surgery is about 1%.

How should I resume activity after hernia surgery?

We recommend that after 2 weeks you slowly resume all of your normal activity and exercises. This will differ from person to person. Start at about 50% of your normal exercise, and progressively increase this over a 1-2 week period of time. We typically recommend starting with aerobic exercises first (running, biking, hiking) before moving to strengthening exercises.

As long as you progress over time and listen to your body on when to slow down, you will not damage the repair.

I have a low grade fever - should I be worried?

A low grade fever is normal after having anesthesia. This is defined as a temperature less than 101F and should resolve within 24 to 48 hours. Please call the office if you have a fever greater than 101F or if it persists longer than 48 hours.

Call the office if you have questions or concerns about your recovery: 617-466-3373

If you are unable to reach your surgeon, please call the hospital at 617-243-6000 and ask the operator to page or connect you to your surgeon